NOAA mobile laboratory measurements of atmospheric emissions from agriculture in northeastern Colorado

Scott Eilerman, Jeff Peischl, Andy Neuman, Tom Ryerson, Anne Perring, Rob Wild, Ken Aikin, Max Holloway

NOAA ESRL Chemical Sciences Division

Cooperative Institute for Research in Environmental Sciences, University of Colorado Boulder

In collaboration with C. Floerchinger, Aerodyne Research, and M. Zondlo, Princeton University









NOAA mobile lab will contribute to our understanding of agricultural impacts on Front Range air quality

- Primary motivation is to quantify and attribute atmospheric emissions of trace carbon and nitrogen species from various point sources in northeastern Colorado, including:
 - Agricultural sites: feedlots, dairies, swine, chickens, sheep
 - Oil and gas wells
 - Landfills and wastewater treatment
- New mobile laboratory enables high resolution measurements with close proximity to the site of interest.

NOAA ICSD IMobile II.aboratory IMeasurements I				
Measured Parameter 2	Method ②	Time Resolution 2	Detection 1 imit 2	G
N ₂ O,3CO,3H ₂ O?	Integrated avity butput spectroscopy 2	132	0.2ppbforIN ₂ O2 0.2ppbfforICO2 100ppmfforIH ₂ O2	[
CO ₂ BandICH ₄ P	Wavelengthscannedscavitysing-downspectroscopy	13?	0.2॒appmvaforaCO₂2 2appbvaforaCH₄2	[i
NH ₃ @nd@H ₂ O@	Cavity Ting-down Spectroscopy 2	variable2	1॒ppbdfordNH₃2 0.05%dfordH₂O2	Œ
$NO_{2}NO_{2}$	Cavity@ing-down@pectroscopy@	13:2	0.1330.0013ppbv2	Ē
WIBS2	Singleparticlefluorescence?	1⊡min⊡	10⁵⊈particles/m³⊡	Œ
GPS@and@meteorology@	Differential IGPS I and I Airmar I Wx I	132	নে	[]






Photo credit: K. Aikin

Preliminary data show clear diurnal, temperature trends

- A clear diurnal cycle and temperature dependence is observed in the NH₃/CH₄ enhancement ratio at several different feedlots.
- The mean enhancement ratio agrees with EPA NEI 2011, but the magnitude of emissions has not yet been quantified.
- Future work will include seasonal variations in emissions, comparison with oil and gas emissions, and a closer examination of the differences between animal types and manure management practices.

